

A TWO-TEMPERATURE MODEL OF THE INTRACLUSTER MEDIUM

MOTOKAZU TAKIZAWA

Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto
606-8502, JAPAN

ABSTRACT

We investigate evolution of the intracluster medium (ICM), considering the relaxation process between the ions and electrons. According to the standard scenario of structure formation, ICM is heated by the shock in the accretion flow to the gravitational potential well of the dark halo. The shock primarily heats the ions because the kinetic energy of an ion entering the shock is larger than that of an electron by the ratio of masses. Then the electrons and ions exchange the energy through coulomb collisions and reach the equilibrium. From simple order estimation we find that the region where the electron temperature is considerably lower than the ion temperature spreads out on a Mpc scale.

We then calculate the ion and electron temperature profiles by combining the adiabatic model of two-temperature plasma by Fox & Loeb (1997) with spherically symmetric N-body and hydrodynamic simulations based on three different cosmological models. It is found that the electron temperature is about a half of the mean temperature at radii ~ 1 Mpc. This could lead to an about 50 % underestimation in the total mass contained within ~ 1 Mpc when the electron temperature profiles are used. The polytropic indices of the electron temperature profiles are $\simeq 1.5$ whereas those of mean temperature $\simeq 1.3$ for $r \geq 1$ Mpc. This result is consistent both with the X-ray observations on electron temperature profiles and with some theoretical and numerical predictions about mean temperature profiles.

Subject headings: galaxies: clustering — galaxies: intergalactic medium — galaxies: X-ray — hydrodynamics

1. INTRODUCTION

Clusters of galaxies (CG) are the largest virialized objects in the universe, which contain collisionless particles, galaxies and dark matter, and a diffuse gas component. The gas

component is called intracluster medium (ICM). The ICM is the plasma with temperature of about 10^8 K, thus emitting X-ray mainly through the thermal bremsstrahlung of the electrons (Sarazin 1988).

Revealing temperature profiles of ICM is an important problem. In theoretical work concerning CG, isothermality of ICM is often assumed. In particular, the isothermal β model (Cavaliere & Fusco-Femiano 1976) is adopted in the conventional mass determination through X-ray observations, in the estimation of the Hubble constant (H_0) through Sunyaev-Zel'dovich (SZ) effect (Sunyaev & Zel'dovich 1972; Birkinshaw, Hughes, & Arnaud 1991), etc. However, deviation from a flat temperature profile becomes important at lower densities in the mass estimation using X-ray data (Evrard, Metzler, & Navarro 1996; Schindler 1996). Also in the H_0 estimation through SZ effect the non-isothermality is one important source of errors (Inagaki, Sugimotohara, & Suto 1995; Yoshikawa, Itoh, & Suto 1998). Therefore, accurate temperature profiles of ICM are required to improve such methods.

On the other hand, temperature maps of ICM provide us with useful information about CG. In merging clusters, characteristic temperature structures are expected to occur through shock heating and adiabatic compression. Some numerical simulations are especially focused on this problem (Schindler & Müller 1993; Ishizaka & Mineshige 1996; Roettiger, Loken, & Burns 1997; Ishizaka 1998; Roettiger, Stone, & Mushotzky 1998). In particular, off-center collisions are investigated by Ricker (1998). Comparing these results with the X-ray observational data, we can guess in what phase the merging clusters are (Fujita et al. 1996; Honda et al. 1996; Churazov et al. 1998; Donnelly et al. 1998; Davis & White 1998).

Recent X-ray observations with ASCA and ROSAT reveal the radial electron temperature distribution of ICM (Markevitch et al. 1996; Markevitch 1996; Markevitch, Sarazin, & Irwin 1996; Markevitch et al. 1997). In some clusters the observed electron temperature gradients correspond to the polytropic index, $\gamma(P \propto \rho^\gamma)$, of $\gamma = 1.5$ or even more (Markevitch 1996). On the other hand, $\gamma \simeq 1.2$ is expected for plasma mean temperature distribution derived by the self similar solution of Bertschinger (1985) and some numerical simulations (Evrard 1990; Katz & White 1993; Navarro, Frenk, & White 1995; Eke, Navarro & Frenk 1997; Takizawa & Mineshige 1998; Bryan & Norman 1998). There is a discrepancy between the observed values and the theoretical expectations.

To explain the discrepancy between the observed electron temperature profiles and the theoretical suggestions on the mean temperature profiles consistently, we construct a model of ICM incorporating properly the relaxation process between the ions and electrons. Fox & Loeb (1997) was the first to investigate the two-temperature nature of ICM. They

construct the electron and ion temperature profiles by combining the self-similar solution by Bertschinger (1985) with the analytical evolutionary model of adiabatic two-temperature plasma which is originally found by Shafranov (1957). Chiéze, Alimi & Teyssier (1998) carried out the 3-D hydrodynamical simulations of two-temperature ICM. However, CGs only in the Einstein de Sitter universe were considered in the previous work concerning the two-temperature model of ICM. Since in a higher density universe CG form at more recent epoch (Richstone, Loeb, & Turner 1992), the previous work is restricted to the case in which a temperature difference is expected to be the largest among reasonable cosmological models. Therefore, it is necessary to study a two-temperature model of ICM in other cosmological models in order to confirm whether the temperature difference is really significant or not. This problem is also related to the dependence of temperature profiles on the cosmological parameters, which is discussed in some numerical simulations (Evrard et al. 1996; Eke et al. 1997). Furthermore, quantification of resultant temperature profiles by using polytropic indices is not fully discussed in the previous work, which is very important to compare the models with X-ray observations properly. Ettori & Fabian (1998) studied two-temperature ICM using simple analytic models and applied their results to the CG A2163. However they neglected the dynamical properties of ICM and assumed that all the ICM in CG is heated at the same time. This assumption is problematic to quantify the temperature profiles because the timescale of shock propagation in CG is comparable to the dynamical timescale and not much shorter than the age of the universe.

For these purpose, we apply the method of Fox & Loeb (1997) to the results of numerical simulations instead of the self-similar solution. Thus we can investigate two-temperature ICM quantitatively in cosmological models other than the Einstein de Sitter model. Studying the dependence of electron temperature profiles on cosmological models is another important purpose.

The rest of this paper is organized as follows. In §2 we estimate the relevant timescales and spatial scales of ICM. In §3 we briefly review the adiabatic model of two-temperature plasma proposed by Fox & Loeb (1997). In §4 we described the method to calculate the ion and electron temperature profiles of ICM by combining a one-temperature ICM model with the adiabatic model by Fox & Loeb (1997). In §5 we describe the adopted numerical methods and initial conditions for simulations of CG. In §6 we present the results. In §7 we summarize the results and discuss their implications.

2. ORDER ESTIMATION

We consider fully ionized plasma which consists of electrons and ions. Only coulomb coupling is considered as the relaxation process. Then two-body relaxation timescale of a x-particle, whose density is n_x and temperature is T_x , is (Spitzer 1962)

$$t_{xx} = \frac{m_x^{1/2}(3kT_x)^{3/2}}{5.71\pi n_x e^4 Z_x^4 \ln \Lambda}, \quad (1)$$

where, m_x is the particle mass, Z_x is the particle charge number, e is the electron charge, k is the Boltzmann constant, and $\ln \Lambda$ is the Coulomb logarithm and approximated to be

$$\ln \Lambda \simeq 37.8 + \ln\left(\frac{T_x}{10^8 \text{K}}\right) - \frac{1}{2} \ln\left(\frac{n_x}{10^{-3} \text{cm}^{-3}}\right) \quad (2)$$

for $T_x > 4 \times 10^5$ K. Therefore, t_{ii} is larger than t_{ee} by a factor of the order of $(m_i/m_e)^{1/2}$.

On the other hand, the equilibrium timescale between ions and electrons is,

$$t_{ei} = \frac{3m_e m_i}{8(2\pi)^{1/2} n_i Z_i^2 e^4 \ln \Lambda} \left(\frac{kT_e}{m_e} + \frac{kT_i}{m_i} \right)^{3/2}. \quad (3)$$

Therefore, t_{ei} is grater than t_{ii} by a factor of $(m_i/m_e)^{1/2}$. In addition, this timescale can be comparable to or longer than the Hubble time in the outer region of CGs,

$$t_{ei} = 2.0 \times 10^8 \text{yr} \left(\frac{\ln \Lambda}{40} \right)^{-1} \left(\frac{n_i}{10^{-3} \text{cm}^{-3}} \right)^{-1} \left(\frac{T_e}{10^8 \text{K}} \right)^{3/2}. \quad (4)$$

Suppose that ICM is heated through the shock in accretion flow (Cavaliere, Menci, & Tozzi 1997). The shock primarily heats ions because the kinetic energy of a particle is proportional to the particle mass. In the post shock region the ions reach thermal equilibrium on a timescale of t_{ii} after they are heated through the shock. Within this time ion temperature is significantly higher than electron one. Eventually thermal energy is transported from the ions to the electrons through the Coulomb collisions between the ions and electrons and T_i becomes comparable to be T_e on timescale of t_{ei} . Note that $t_{ei} > t_{ee}, t_{ii}$.

Under such circumstance, the radial length, r_{tt} , over which the electron temperature is significantly lower than the ion temperature can be estimated as follows. Denote the propagation speed of the shock front by v_{shock} . Then, we have,

$$r_{tt} \simeq t_{ei} v_{\text{shock}}. \quad (5)$$

Using the strong shock approximation and neglecting the post-shock gas velocity compared with v_{shock} , we find

$$v_{\text{shock}} \simeq \frac{1}{3} v_{\text{infall}}, \quad (6)$$

where v_{infall} is the infalling velocity of the gas, which is related to the post-shock gas temperature, T . The kinetic energy of infalling gas is nearly equal to the thermal energy of the post shock gas; namely

$$\frac{1}{2}m_p v_{\text{infall}}^2 \simeq k_B T. \quad (7)$$

Therefore, using the equations (4), (5), (6), and (7), we derive

$$r_{\text{tt}} \simeq 1.1 \times 10^{-1} \text{Mpc} \left(\frac{\ln \Lambda}{40} \right)^{-1} \left(\frac{n_i}{10^{-3} \text{cm}^{-3}} \right)^{-1} \left(\frac{T_e}{10^8 \text{K}} \right)^{-2}. \quad (8)$$

Importantly this is smaller by about one order of magnitude than the spatial scale of CGs. Note that the higher temperature is, the wider becomes the region where the temperature difference between ions and electrons is significant.

Another important timescale related to ICM is the radiative cooling timescale, t_c . Since thermal bremsstrahlung is the dominant cooling process in ICM, we have

$$t_c = 8.5 \times 10^{10} \text{yr} \left(\frac{n_e}{10^{-3} \text{cm}^{-3}} \right)^{-1} \left(\frac{T_e}{10^8 \text{K}} \right)^{1/2}. \quad (9)$$

From equations (4) and (9) we find,

$$\frac{t_{\text{ei}}}{t_c} = 2.3 \times 10^{-3} \left(\frac{\ln \Lambda}{40} \right)^{-1} \left(\frac{T_e}{10^8 \text{K}} \right). \quad (10)$$

Hence t_c is always longer than t_{ei} in the typical ICM, and thus we can safely neglect cooling effects as far as we are concerned with the overall cluster structure.

On the other hand, the age of CG, t_{age} , is of the order of 10^9 or 10^{10} yr. Therefore, as long as heating from galaxies can be neglected, we can divide intra-cluster space into three regions according to the magnitudes of these three timescales, t_{ei} , t_c , and t_{age} as follows:

1. The central higher density region where $t_{\text{ei}} < t_c < t_{\text{age}}$. Radiative cooling is important but ICM can be regarded as one-temperature fluid. This situation corresponds to the so-cold 'cooling flow'.
2. The middle region where $t_{\text{ei}} < t_{\text{age}} < t_c$. We can regard ICM as adiabatic one-temperature fluid.
3. The outer lower density region where $t_{\text{age}} < t_{\text{ei}} < t_c$. Radiative cooling can be negligible and the electron temperature is considerably lower than the ion temperature.

3. EVOLUTION OF ADIABATIC TWO-TEMPERATURE PLASMA

The formulation here is based on Fox & Loeb(1997). We neglect thermal conduction, which is the case if tangled magnetic fields suppress conduction. The Lagrangean time evolution of the electron temperature, T_e , and the mean temperature, $\bar{T} = (n_e T_e + n_i T_i)/(n_e + n_i)$, in the adiabatic fluid element is,

$$\frac{dT_e}{dt} = \frac{T_i - T_e}{t_{ei}} + (\gamma - 1) \frac{T_e}{n} \frac{dn}{dt}, \quad (11)$$

$$\frac{d\bar{T}}{dt} = (\gamma - 1) \frac{\bar{T}}{n} \frac{dn}{dt}, \quad (12)$$

where T_i is the ion temperature, n is the gas density, and $\gamma = 5/3$, is the ratio of specific heat. Introducing the temperatures normalized by $\bar{T}$, $\tilde{T}_e \equiv (T_e/\bar{T})$ and $\tilde{T}_i \equiv (T_i/\bar{T})$, we find,

$$\frac{d\tilde{T}_e}{dt} = \frac{\tilde{T}_i - \tilde{T}_e}{t_{ei}}. \quad (13)$$

Note that t_{ei} is proportional to $\tilde{T}_e^{3/2} \ln \Lambda$, since in the post-shock region the gas behaves adiabatically ($\bar{T} \propto n^{2/3}$). Thus $t_{2s} \equiv t_{ei}(t) \tilde{T}_e(t)^{-3/2}$ is constant in time, if we neglect the small change due to the Coulomb logarithm. Now equation (13) becomes

$$\frac{d\tilde{T}_e}{dt} = \frac{1}{t_{2s}} \left(\frac{n_i + n_e}{n_i} \right) (1 - \tilde{T}_e) \tilde{T}_e^{-3/2}. \quad (14)$$

Since ICM is almost perfectly ionized, the ratio of $(n_i + n_e)/n_i$ can be regarded as constant. Thus equation (14) can be integrated analytically. If we assume the pre-shock $\tilde{T}_e$ equal to zero, the solution for the fluid element which has passed the shock front at $t = t_s$ is,

$$t - t_s = t_{2s} \left(\frac{n_i}{n_i + n_e} \right) \left[\ln \left(\frac{1 + \sqrt{\tilde{T}_e}}{1 - \sqrt{\tilde{T}_e}} \right) - 2\sqrt{\tilde{T}_e} \left(1 + \frac{\tilde{T}_e}{3} \right) \right]. \quad (15)$$

Thus we can obtain the Lagrangean time evolution of $\tilde{T}_e$ of the fluid element after the passage through the shock by solving equation (15).

4. NUMERICAL METHOD

Using equation (15) we can construct the temperature profiles at $t = t_0$, $T_e(r, t_0)$ and $T_i(r, t_0)$, as follows. First, we give the velocity field of the gas, $v(r, t)$, the shock radius, $r_{\text{shock}}(t)$, and the equilibrium timescale of the fluid element at the passage through the

shock front, $t_{2s}[r_{\text{shock}}(t)]$, using some analytical models or numerical simulations presented later. Then we can calculate the Lagrangean path of the fluid element which passed the shock surface at $t = t_s$, $R(t; t_s)$, by solving the ordinary differential equation,

$$\frac{dR}{dt} = v(R(t; t_s), t), \quad (16)$$

with the appropriate initial condition,

$$R(t_s; t_s) = r_{\text{shock}}(t_s). \quad (17)$$

Integrating differential equation (16) for various t_s , we obtain $r_0 = R(t_0; t_s)$, the radius which the fluid element that passed the shock at $t = t_s$ resides at $t = t_0$, as a function of t_s . Thus we can regard $t_0 - t_s$ as a function of the radius,

$$t_0 - t_s = f(r; t_0). \quad (18)$$

Using the equations (15) and (18), and the model of $t_{2s}[r_{\text{shock}}(t_s)]$, we can solve $\tilde{T}_e$ at $t = t_0$ as a function of r . Finally, we obtain $T_e(r)$ and $T_i(r)$ using $\tilde{T}_e(r)$ and the model of $\bar{T}(r)$.

5. THE SIMULATIONS

5.1. Numerical Method

To give $v(r, t)$, $r_{\text{shock}}(t)$, etc, we have performed numerical simulations of a spherically symmetric CG. For dark matter (DM), we use the shell model (Hénon 1964). We set the number of shells, N , equal to 5000. As for gas on the other hand, we use 1-dimensional, spherically symmetric, total variation diminishing (TVD) code with minmod limiter (Hirsch 1990). Note that TVD code is one of the most powerful tools to treat shocks. One mesh spacing corresponds to $\Delta r = 0.005/(1+z)$ Mpc. We assume that the gas is ideal, with $\gamma = 5/3$. As to the boundary conditions, the inner edge is assumed to be a perfectly reflecting point. The outer edge is assumed to be a perfectly transmitting surface. The basic equations and the numerical method used here are fully described in §2 of Takizawa & Mineshige (1998).

5.2. Models and Initial Conditions

In this paper, all of the calculations are carried out from $z_{\text{ini}} = 10$ to the present time ($z_0 = 0$). The cosmological models which we adopt are $(\Omega_0, \Lambda_0) = (1.0, 0.0)$ (ED),

$(\Omega_0, \Lambda_0) = (0.2, 0.0)$ (OP), and $(\Omega_0, \Lambda_0) = (0.2, 0.8)$ (FL). The Hubble constant, H_0 is set to be $H_0 = 100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ in the transformations of length and time coordinates. We set $\Omega_b = 0.012 h^{-2}$ in all models taken from the nucleosynthesis determination (Walker et al. 1991). Note that $\Omega_{\text{DM}} = \Omega_0 - \Omega_b$.

We make initial density profiles in the same manner as Peebles (1982). At first we prepare N concentric shells with a constant density equal to Ω_{DM} at $z_{\text{ini}} = 10$. Then, a density fluctuation is introduced by perturbing the radius and velocity of each shell following

$$r_i = r_i^{(0)} \left\{ 1 - \frac{1}{3} \bar{\delta}[r_i^{(0)}] \right\}, \quad (19)$$

$$u_i = H(z_{\text{ini}}) r_i^{(0)} \left\{ 1 - \frac{1}{3} (1 + \Omega^{0.6}) \bar{\delta}[r_i^{(0)}] \right\}, \quad (20)$$

where $r_i^{(0)}$ is the unperturbed coordinate, $\bar{\delta}(r)$ represents the mean density fluctuation inside r , which is derived from the density fluctuation field (specified below) within r , and $H(z_{\text{ini}})$ is the Hubble constant at $z = z_{\text{ini}}$. Here, we used the Zel'dovich approximation (Zel'dovich 1970) and the approximation of $d \log D_1 / d \log a \simeq \Omega^{0.6}$, where D_1 is the linear perturbation growth rate for the growing mode and a is the scale factor (see Suto 1993).

The initial conditions of the density fluctuation field, $\delta(r)$, are generated by applying the Hoffman-Ribak method (Hoffman & Ribak 1991; van de Weygaert & Bertschinger 1996) to spherical systems (see Appendix). We constrained in such a way that there exist initially density enhancements on 1 Mpc scale whose amplitudes correspond to 3σ level in the CDM power spectrum. The normalization is $\sigma_8 = 1$ in each cosmological model, which was obtained from the observation of the nearby galaxy distribution (see Suto 1993).

The initial conditions of gas are set as follows. At first, gas density was everywhere taken to be the mean baryon density of the universe at $z = 10$, and the temperature of the gas ($T_{\text{gas},i}$) was constant everywhere; $T_{\text{gas},i} = 10^7 \text{ K}$ in all models. We then add the adiabatic fluctuation in such a way that the ratio of the DM density and the gas density remains the same. Note that after the perturbation is added, the gas temperature distribution becomes nonuniform, accordingly. Moreover, note that temperature of the infalling gas at $z \sim 1$ is sufficiently lower than virial temperature, since the gas expands adiabatically following the cosmological expansion.

6. RESULTS

The overall evolution of the simulated CG is essentially the same as that presented by Takizawa & Mineshige (1998). Thus we concentrate on showing the results of the temperature profiles, mass estimation, and so on.

6.1. Temperature Profiles

Figure 1 shows radial distribution the $T_e(r)$ (solid line), $T_i(r)$ (dotted line), and $\bar{T}(r)$ (short dashed line) of Model ED at $z = 0$. At $r < 0.6\text{Mpc}$ T_e is very close to T_i , while, at $r > 0.6\text{Mpc}$ the electron temperature is considerably lower than the ion temperature and the discrepancy gets increased outward. At $r \simeq 1\text{Mpc}$, the electron temperature is only a half of the mean temperature. Small scale fluctuations in the temperature profiles are due to the sound wave propagation in ICM (Takizawa & Mineshige 1988).

In real X-ray observations what is actually obtained is an emissivity-weighted, line-of-sight projected electron temperature map, which is displayed in Figure (2) for model ED. In this figure, we assumed the spatial resolution to be 0.25 Mpc, which corresponds to about 6' for an object located at $z = 0.05$, and the error of the temperature measurement to be 20 %. Also in this map we can clearly see that the electron temperature decreases outward.

To estimate the temperature gradients both for T_e and $\bar{T}$, we measure the polytropic indexes, γ_p , in the usual way. We fit the density profile by the β -model and the temperature profile by the polytropic model as follows,

$$n(r) = n_0 \left[1 + \left(\frac{r}{r_c} \right)^2 \right]^{-3\beta/2}, \quad (21)$$

$$T(r) \propto n(r)^{\gamma_p - 1} \quad (22)$$

where n_0 , r_c , and β is the fitting parameters of the β -model. We fit the resultant density profile only inside the shock front. The data are fitted by chi-square fitting. We assume that the variance in any quantity is proportional to its square ($\delta f/f = \text{const.}$) because our main purpose is not to simulate observations with a specific instrument but to obtain the intrinsic profiles of our calculated results. The results of the fitting are summarized in Table 1 and the corresponding polytropic indices are listed in Table 2. From Table 1 we find that the β values are somewhat bigger than those typically observed. On the other hand, our results are well coincident with the self-similar solution by Bertschinger (1985). Thus the bigger β values are probably due to the assumption of spherical symmetry and neglect

of angular momentum of gas. If the initial gas temperature is higher, which corresponds to the case that reheating of the ICM, e.g., by protogalaxies, is substantial, β values can be smaller (Metzler & Evrard 1994; Takizawa & Mineshige 1988). From Table 2 we find that the polytropic indices of the electron temperature profiles are $\gamma_p \sim 1.5$, which are systematically larger than those of the mean temperature profiles, $\gamma_p \sim 1.3$.

The specific entropy profiles derived from the electron temperature [$S_e \propto \ln(T_e/\rho^{\gamma-1})$], and from the mean temperature [$\bar{S} \propto \ln(\bar{T}/\rho^{\gamma-1})$], are shown in Figure 3 by the solid and dotted lines, respectively. Entropy is normalized to be zero at the inner boundary. The latter $\bar{S}$ rises outward, which is characteristic of the ICM heated through the shock (Evrard 1990; Takizawa & Mineshige 1998). The former S_e , in contrast, rises outward as the case only in the inner region ($r < 0.6\text{Mpc}$), stays nearly constant in the middle region, and falls outward in the outer region.

6.2. Dependence of Fitting Parameters on Size of Fitting

We fit the calculated density profile inside the shock front by equation (21) and listed the fitting results in Tables 1 and 2. However, it is possible that the fitting results can be influenced by the position of the outer edge of the region used for the fitting. In real X-ray observations the outer edge of the X-ray emitting region is perhaps inside the shock front. To assess this effect we fit the density and temperature profiles inside various radii for Model ED and list the dependence of the fitting results related to the density profile (n_0 , r_c , and β) on the outer-edge radius (r_{out}) in Figure 4. It is found that these parameters are insensitive to the outer radius as long as $r_{\text{out}} > 0.8\text{ Mpc}$. In density profiles, therefore, we can safely neglect the influence of the outer edge. This fact is actually expected because of the self-similar nature of the gas density profile (Bertschinger 1985; Takizawa & Mineshige 1998).

On the other hand, the dependence of γ_p on r_{out} is rather different. Figure 5 shows that γ_p derived from T_e (by closed squares) monotonically increase outward whereas γ_p derived from $\bar{T}$ (by open square) does not exhibit systematic changes. When we calculate a polytropic index from X-ray observational data, thus, the resultant value could be subjected to large errors arising from the finite detection limit, background noise, and so on.

The results of model OP and FL are essentially the same as those of model ED. Although there is a simple self-similar solution only for model ED, self-similar nature is also expected both in OP and FL (Takizawa & Mineshige 1988).

6.3. Mass Estimation

Since the observed electron temperature significantly deviates from the mean temperature that determines the dynamics of the system, the total mass of CG is probably underestimated if hydrostatic equilibrium is calculated based on the electron temperature map. When ICM is assumed to be isothermal we usually use the emissivity-weighted mean temperature, which is more like the temperature in the central high-density region. Thus, the underestimation of the mass is practically negligible. When the electron temperature profile is used, conversely, the mass can be seriously underestimated. Hence the mass derived from the assumption of hydrostatic equilibrium, $M_{\text{hydro}}(r)$, is,

$$M_{\text{hydro}}(r) = \frac{3kT_0\beta\gamma_p}{\mu m_p G} r \frac{(r/r_c)^2}{[1 + (r/r_c)^2]^{1+3\beta(\gamma_p-1)/2}}. \quad (23)$$

Figure 6 depicts the ratio of M_{hydro} to the actual mass as a function of radius. The solid line represents the ratio calculated based on $T_e(r)$ and the dotted line represents that based on $\bar{T}(r)$. The mass derived from T_e is underestimated by almost 50 % because of the lower electron temperature. Note that the mass derived from $\bar{T}$ is also underestimated by about 10 % due to the bulk motion of the gas.

In the central region M_{hydro} based on T_e and $\bar{T}$ are both larger than the actual mass. This is due to the fitting errors of temperature profiles. Since the core radii are different between the density and temperature profiles, polytropic model cannot describe our results well in this region.

7. SUMMARY AND DISCUSSION

We constructed the models of ICM, incorporating the relaxation process between the ions and electrons. From the simple order estimation, we find that the electron temperature is well below the ion temperature in the outer region of CG and that such a lower T_e region spreads over a Mpc scale in typical CG. In addition, the hotter CG is, the wider becomes the two-temperature region. Comparing three relevant timescales in ICM (the age of CG, radiative cooling, and equilibrium timescales between ions and electrons), we can divide ICM into three regions; from the center outward, the cooling dominant, one-temperature region, adiabatic, one-temperature region, and adiabatic, two-temperature region.

We calculate the temperature profiles of two-temperature ICM combining the spherically symmetric, N-body and hydrodynamic simulations for three different cosmological models with the adiabatic two-temperature plasma model by Fox & Loeb (1997). While the

polytropic indices of the mean temperature profiles are $\simeq 1.3$, those of the electron temperature profiles are $\simeq 1.5$. As a consequence, the specific entropy profiles derived from the electron temperature are rather flat.

We examine the dependence of the fitting parameters on the radius of the outer edge of the region used for the fitting when the density profile is fitted by the β -model and the temperature profile by the polytropic model. The fitting results of the density profile and mean temperature profile is insensitive to the outer edge. On the other hand, the polytropic index derived from the electron temperature rises as the outer-edge radius increases.

The total mass of CG is underestimated about 50 % when we use the electron temperature profile though the underestimation is negligible when we assume that ICM is isothermal and adopt emission-weighted mean temperature as the temperature of ICM.

We confirm that temperature difference between ions and electrons in ICM is substantial in cosmological models other than the Einstein de Sitter model. The polytropic indices of electron temperature profiles are insensitive to the cosmological models in the range of our calculations. Note that the baryon density of these models is set to be the same. If the baryon fraction is set to be constant in each model, the result will probably changes because the equilibrium timescale is sensitive to the baryon density.

In general our resultant electron-temperature profiles tend to have steeper gradients than those of Ettori & Fabian (1998). They assumed that all the ICM in CG is heated at the same time and that ICM is in hydrostatic equilibrium. On the other hand, in our calculations the ICM in the outer region is heated more recently than that in the inner region because the shocks propagate at finite speed. Furthermore, gradual radial infall persists inside the shock fronts and makes the two-temperature region slightly compress inward. (Takizawa & Mineshige 1998). Thus the two-temperature region of our results becomes wider than those of Ettori & Fabian (1988).

Although only spherical accretion is considered in this paper, there may also arise asymmetric merging of comparable clumps in reality. It is possible that shocks occurring in merging events will also generate the temperature difference between ions and electrons. In this case it is believed that a bow shock with an arc shape is formed just between the centers of two substructures (Schindler & Müller 1993; Ishizaka & Mineshige 1996; Roettiger et al. 1997; Ishizaka 1998). In the post-shock region the energy is transported from ions to electrons but that timescale changes along the shock front since the timescale is sensitive to the density of ICM. Therefore, if we consider the temperature difference between ions and electrons, the location of the observed hot gas region is probably shifted and the shape is more deformed in comparison with the results of the former simulations.

We will investigate this issue as a future work.

We consider only the classical coulomb coupling as the relaxation process between ions and electrons. It is possible, however, in ICM more efficient relaxation processes may be effective (McKee & Cowie 1977; Pistinner, Levinson, & Eichler 1996). In this case the equilibrium timescale can be shorter than the value given by equation (4). Therefore, the temperature difference between ions and electrons can be less and polytropic indices can be smaller than our results. If magnetic field exists in ICM, it is possible that electrons are also significantly heated in shocks by MHD instabilities. Also in this case the temperature difference between ions and electrons can be less.

We neglect the heating process from the galaxies to ICM. If thermalized hot gas is injected to ICM from the galaxies, temperature difference could be less. Furthermore, in this case the preheating of the ICM influences the density profile and mean temperature profile (Metzler & Evrard 1994; Navarro et al. 1995; Takizawa & Mineshige 1998). Since heavy elements like iron are detected in ICM at redshift up to $z \simeq 1$ (Hattori et al. 1997), this effect should be considered to construct a more realistic model. In such a study the adiabatic model like Fox & Loeb (1997) cannot be adopted. Thus non-adiabatic models or fully two-fluid simulations like Chiéze et al. (1998) are required.

The author would like to thank S. Mineshige for valuable comments. This work is supported by a Grant-in-Aid from the Ministry of Education, Science, Sports and Culture of Japan (6179) and Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

A. THE HOFFMAN-RIBAK METHOD FOR SPHERICAL SYSTEMS

We consider a random homogeneous and isotropic Gaussian field f with zero mean which is defined by its power spectrum $P(k)$. When f is subjected to linear constraints, $g = Cf$, then the constrained Gaussian field f is realized as follows (Hoffman & Ribak 1991),

$$f = \tilde{f} + MC^\dagger Q^{-1}(g - C\tilde{f}), \quad (\text{A1})$$

where $\tilde{f}$ is an unconstrained random Gaussian field whose power spectrum is $P(k)$, M is the two-point correlation matrix obtained from $P(k)$, and $Q = CMC^\dagger$.

In general three dimensional case, $\tilde{f}$ is described by its Fourier components $\tilde{f}_{\mathbf{k}}$,

$$\tilde{f}(\mathbf{r}) = \frac{1}{(2\pi)^3} \int \tilde{f}_{\mathbf{k}} \exp(i\mathbf{k}\mathbf{r}) d\mathbf{k}. \quad (\text{A2})$$

However, to generating the initial conditions which can be used for our spherical symmetric simulations, we should construct $\tilde{f}$ as a function of only r instead of $\mathbf{r}$. Since $\tilde{f}$ is isotropic $\tilde{f}(r)$ is obtained as follows. In spherical systems it is convenient to expand $\tilde{f}(\mathbf{r})$ in spherical harmonics times spherical Bessel functions j_l (Binney & Quinn 1991). A plain wave is expanded in these functions as follows,

$$\exp(i\mathbf{k}\mathbf{r}) = 4\pi \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} i^l j_l(kr) Y_l^{m*}(\theta_k, \phi_k) Y_l^m(\theta_r, \phi_r). \quad (\text{A3})$$

Integrating this for the tangential components we find,

$$\int \exp(i\mathbf{k}\mathbf{r}) d\mathbf{k} = 4\pi \int j_0(kr) k^2 dk. \quad (\text{A4})$$

We can assume that $\tilde{f}_{\mathbf{k}} = \tilde{f}_k$ because of isotropy of $\tilde{f}$. Thus we obtain,

$$\tilde{f}(r) = \frac{1}{2\pi^2} \int \tilde{f}_k \frac{\sin kr}{kr} k^2 dk. \quad (\text{A5})$$

Model	n_0 (10^{-3}cm^{-3})	r_c (Mpc)	β
ED	1.16	0.067	0.793
OP	3.64	0.104	0.815
FL	1.45	0.155	0.903

Table 1: Density profiles of gas at $z = 0$.

Model	γ_p of T_e	γ_p of $\bar{T}$
ED	1.55	1.35
OP	1.44	1.34
FL	1.51	1.37

Table 2: Polytropic indices of T_e and $\bar{T}$ at $z = 0$.

REFERENCES

- Bertschinger, E. 1985, ApJS, 58, 39
- Binney, J., & Quinn, T. 1991, MNRAS, 249, 678
- Birkinshaw, M., Hughes, J. P., & Arnaud, K. A. 1991, ApJ, 379, 466
- Bryan, G. L., & Norman, M. L. 1997, in Computational Astrophysics, Proc. 12th Kingston Conference, eds. D Clarke & M. West (PASP), astro-ph/9710186
- Cavaliere, A., & Fusco-Femiano, R. 1976, A&A, 49, 137
- Cavaliere, A., Menci, N., & Tozzi, P. 1997, ApJ, 484, L21
- Chiéze, J.-P., Alimi, J.-M., & Teyssier, R. 1998, ApJ, 495, 630
- Churazov, E., Gilfanov, M., Forman, W., & Jones, C. 1998, ApJ, submitted, astro-ph/9802166
- Davis, D. S. & White III, R. E. 1998, ApJ, 492, 57
- Donnelly, R. H., Markevitch, M., Forman, W., Jones, C., David, L. P., Churazov, E., & Gilfanov, M. 1998, ApJ, 500, 138
- Eke, V. R., Navarro, J. F., & Frenk, C. S. 1997, ApJ, submitted, astro-ph/9708070
- Ettori, S. & Fabian, A. C. 1998, MNRAS, 293, L33
- Evrard, A. E. 1990, ApJ, 363, 349
- Evrard, A. E., Metzler, C. A., & Navarro, J. F. 1996, ApJ, 469, 494
- Fox, D. C., & Loeb, A. 1997, ApJ, 491, 459
- Fujita, Y., Koyama, K., Tsuru, T., & Matsumoto, H. 1996, PASJ, 48, 191
- Hattori, M., Ikebe, Y., Asaoka, I., Takeshima, Y., Boehringer, H., Mihara, T., Neumann, D. M., Schindler, S., Tsuru, T. & Tamura, T. 1997, Nature, 388, 146
- Hénon, M. 1964, Ann. Astrophys., 27, 83
- Hirsch, C. 1990, Numerical Computation of Internal and External Flows Vol.2: Computational Methods for Inviscid and Viscous Flows (New York: John Wiley & Sons)
- Hoffman, Y., & Ribak, E. 1991, ApJ, 380, L5
- Honda, H., Hirayama, M., Watanabe, M., Kunieda, H., Tawara, Y., Yamashita, K., Ohashi, T., Hughs, J. P., & Henry, J. P. 1996, ApJ, 473, L71
- Inagaki, Y., Sugimotohara, T., & Suto, Y. 1995, PASJ, 54, 411
- Ishizaka, C., & Mineshige, S. 1996, PASJ, 48, L37

- Ishizaka, C., 1998 A & SpS, in press
- Katz, N., & White, S. D. M. 1993, ApJ, 412, 455
- McKee, C. F., & Cowie, L., L. 1977, ApJ, 215, 213
- Markevitch, M., Mushotzky, R., Inoue, H., Yamashita, K., Furuzawa, A., & Tawara, Y. 1996, 456, 437
- Markevitch, M. 1996, ApJ, 465, L1
- Markevitch, M., Sarazin, C. L., & Irwin, J. A. 1996, ApJ, 472, L17
- Markevitch, M., Forman, W. R., Sarazin, C. L., & Vikhlinin, A. 1997, ApJ, submitted, astro-ph/9711289
- Metzler, C. A., & Evrard, A. E. 1994, ApJ, 437, 564
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1995, MNRAS, 275, 720
- Peebles, P. J. E. 1982, ApJ, 257, 438
- Pistinner, S., Levinson, A., & Eichler, D. 1996, ApJ, 467, 162
- Richstone, D., Loeb, A., & Turner, E. L. 1992, ApJ, 393, 477
- Ricker, P., M. 1998, ApJ, 496, 670
- Roettiger, K., Loken, C., & Burns, J. B. 1997 ApJS, 109, 307
- Roettiger, K., Stone, J. M., & Mushotzky, R. F. 1998, ApJ, 493, 62
- Sarazin, C. L. 1988, X-ray Emission from Clusters of Galaxies (Cambridge: Cambridge University Press)
- Shafranov, V. D. 1957, Soviet Phys. JETP, 5, 1183
- Schindler, S., & Müller, E. 1993, A&A, 272, 137
- Schindler, S. 1996, A&A, 305, 756
- Spitzer, L., Jr. 1962, Physics of Fully Ionized Gases (New York: Wiley)
- Sunyaev, R. A., & Zel'dovich, Ya. B. 1972, Comm. Astrophys. Sp. Phys. 4, 173
- Suto, Y. 1993, Prog. Theor. Phys., 90, 1173
- Takizawa, M., & Mineshige, S. 1998, ApJ, 499, 82
- van de Weygaert, R., & Bertschinger, E. 1996, MNRAS, 281, 84
- Walker, T. P., Steigman, G., Schramm, D. N., Olive, K. A., & Kang, H. 1991, ApJ, 376, 51
- Yoshikawa, K., Itoh, M., & Suto, Y. 1998, PASJ, 50, 203

Zel'dovich, Ya. B. 1970, A&A, 5, 84

Fig. 1.— The profiles of the T_e (solid line), T_i (dotted line), and $\bar{T}$ (short dashed line) of Model ED at $z = 0$. At $r < 0.6$ Mpc $T_e \sim T_i$, whereas at $r > 0.6$ Mpc T_e is considerably lower than T_i and the discrepancy gets enhanced outward. At $r \simeq 1$ Mpc, $T_e \sim \bar{T}/2$. Small scale fluctuations in the temperature profiles are due to the sound wave propagation in ICM.

Fig. 2.— Emissivity weighted, line-of-sight projected electron temperature profile of Model ED at $z = 0$. Here we assumed the spatial resolution to be 0.25 Mpc, which corresponds to about 6' for an object located at $z = 0.05$, and the error of the temperature measurement to be 20 %. Clearly the electron temperature decreases outward.

Fig. 3.— The specific entropy profiles derived from the electron temperature [$S_e \propto \ln(T_e/\rho^{\gamma-1})$], by the solid line and from the mean temperature [$\bar{S} \propto \ln(\bar{T}/\rho^{\gamma-1})$], by the dotted line, respectively. Entropy is normalized to be zero at the inner boundary.

Fig. 4.— The dependence of the parameters related to the density profile (n_0 , r_c , and β) on r_{out} , the radius of the outer edge of the region used for fitting. It is found that these parameters are insensitive to the outer edge radius as long as $r_{\text{out}} > 0.8$.

Fig. 5.— The dependence of γ_p on r_{out} . We find that γ_p obtained from T_e (closed squares) increases as r_{out} increases, whereas γ_p obtained from $\bar{T}$ (open squares) is insensitive to r_{out} .

Fig. 6.— The ratio of M_{hydro} to the actual mass. The solid line is the result based on $T_e(r)$ and the dotted line is that based on $\bar{T}(r)$. The mass derived from T_e is underestimated by almost 50 % because of the lower electron temperature. Note that the mass derived from $\bar{T}$ is also underestimated by about 10 % due to the bulk motion of the gas











